

L2 ANSWER 2 OF 3 CAPLUS COPYRIGHT 2003 ACS on STN  
 AN 1999:699081 CAPLUS  
 DN 131:314219  
 TI Compositions containing **Echinacea** extracts and antiallergic agents for common cold  
 IN Asano, Toshinori; Sakata, Yasuko  
 PA Taisho Pharmaceutical Co., Ltd., Japan  
 SO Jpn. Kokai Tokkyo Koho, 4 pp.  
 CODEN: JKXXAF  
 DT Patent  
 LA Japanese  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 11302189	A2	19991102	JP 1998-110797	19980421
PRAI	JP 1998-110797		19980421		

AB Compns. contg. **Echinacea** exts. and antiallergic agents selected from carbinoxamine maleate, chlorpheniramine maleate, brompheniramine maleate, ketotifen fumarate, epinastine-HCl and mequitazine for common cold are claimed. Tablets were formulated contg. acetaminophen 900, codeine phosphate 18, methylepherin HCl 60, carbinoxamine maleate 12, **Echinacea** purpurea exts. 24, **guaifenesin** 125, anhyd. caffeine 50, lactose 275, low-substitution hydroxypropylcellulose 275, magnesium stearate 32 and hardened castor oil 29 g.

L2 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2003 ACS on STN  
 AN 1998:728292 CAPLUS  
 DN 130:17232  
 TI Common cold medicines containing **Echinacea** and antipyretic analgesics  
 IN Sakata, Yasuko; Okuhira, Ichiro; Sumida, Kenji  
 PA Taisho Pharmaceutical Co., Ltd., Japan  
 SO Jpn. Kokai Tokkyo Koho, 5 pp.  
 CODEN: JKXXAF  
 DT Patent  
 LA Japanese  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 10298088	A2	19981110	JP 1997-107849	19970424
PRAI	JP 1997-107849		19970424		

AB Oral compns. contg. **Echinacea** ext. and antipyretic analgesics at the wt. ratio of 1 to (0.1-50) do not have bitterness, therefore are effective as a remedy for common cold with patients' compliance. A tablet (300 mg each) was obtained from a mixt. contg. acetaminophen 900, dihydrocodeine phosphate 24, methylephedrine.cntdot.HCl 60, chlorpheniramine maleate 6, **Echinacea** ext. 24, **guaifenesin** 125, anhyd. caffeine 50, lactose 275, hydroxypropyl cellulose 275, Mg stearate 35, hydrogenated castor oils 26 g.

=>

L4 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2003 ACS on STN  
 AN 2001:114960 CAPLUS  
 DN 134:168363  
 TI **Echinacea** binder for pharmaceutical compositions  
 IN First, Sigal; Yamin, Rina  
 PA Cts Chemical Industries Ltd., Israel  
 SO PCT Int. Appl., 14 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001010415	A1	20010215	WO 2000-IL412	20000713
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
	EP 1200069	A1	20020502	EP 2000-944197	20000713
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL			
PRAI	IL 1999-131317	A	19990809		
	WO 2000-IL412	W	20000713		
AB	Pharmaceutical compns., which contain a binder that comprises a binding-effective amt. of <b>Echinacea</b> prepn. are described. Paracetamol tablets were prepd. with <b>Echinacea</b> as a single binder.				
RE.CNT 4	THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT				

L13 ANSWER 15 OF 30 CAPLUS COPYRIGHT 2003 ACS on STN  
 AN 1998:776655 CAPLUS  
 DN 130:29238  
 TI Pharmaceutical compositions containing NSAIDS  
 IN Barrett, David Michael; Jones, Huw Lyn; Jones, Idwal; Smith, Carl Simon  
 PA The Boots Company PLC, UK  
 SO PCT Int. Appl., 25 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9852540	A1	19981126	WO 1998-EP3179	19980522
	W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			

	AU 9881079	A1	19981211	AU 1998-81079	19980522
PRAI	GB 1997-10505		19970522		
	GB 1997-10527		19970522		
	GB 1997-10544		19970522		
	WO 1998-EP3179		19980522		

AB The present invention relates to the use of an NSAID selected from ibuprofen, naproxen, ketoprofen, diclofenac, piroxicam and indomethacin in the treatment of the symptoms of cold and flu particularly sore throat. The method consists of administration to a patient of a pharmaceutical compn. in the form of a masticable or suckable solid dosage form or a liq. or a spray contg. a therapeutically effective amt. of the NSAID which releases the NSAID in the oral cavity so as to deliver the NSAID to the surface of the sore throat. The compn. may also contain (a) therapeutically effective amt. of 1 or more active ingredients selected from an antihistamine, a cough suppressant, a decongestant, an expectorant, a muscle relaxant, a centrally acting analgesic, a local anesthetic, an antibacterial compd., an antiviral compd., an antibiotic compd., an antifungal compd., minerals and vitamins and/or (b) a burn-masking amt. of an agent which has a warming effect on the mucosa of the throat. Thus, a lozenge contained CaCO<sub>3</sub> 7.5, PVP 1.43, aerosil 0.036, Mg stearate 0.18, isomalt 1885, lycasin 440 mg, ketoprofen q.v. (quantum vis) and flavoring q.v.

RE.CNT 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FO

L10 ANSWER 50 OF 54 CAPLUS COPYRIGHT 2003 ACS on STN  
 AN 1991:415595 CAPLUS  
 DN 115:15595  
 TI Sustained-release pharmaceutical preparation having coated drug  
 microparticles  
 IN Eichel, Herman J.; Massmann, Brent D.  
 PA Kinaform Technology, Inc., USA  
 SO Eur. Pat. Appl., 16 pp.  
 CODEN: EPXXDW  
 DT Patent  
 LA English  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 391518	A2	19901010	EP 1990-301065	19900201
	EP 391518	A3	19910508		
	EP 391518	B1	19930929		
	R: CH, DE, FR, GB, IT, LI, NL, SE				
	US 5026559	A	19910625	US 1989-332154	19890403
	HU 74088	A2	19961128	HU 1990-583	19900130
	HU 214576	B	19980428		
	AU 9050792	A1	19901004	AU 1990-50792	19900306
	AU 622526	B2	19920409		
	JP 02289512	A2	19901129	JP 1990-73152	19900322
	DD 299946	A5	19920514	DD 1990-339325	19900402
PRAI	US 1989-332154		19890403		

AB A sustained-release pharmaceutical prepn. comprises an admixt. of  
 uncoated, single-walled coated, and multi-walled coated microparticles of  
 a drug. The microparticle structure preferably has a core drug, an inner  
 wall microencapsular enteric coating (e.g. polymethacrylic acid/acrylic  
 acid copolymer, cellulose acetate phthalate, etc.), a solid acid (e.g.  
 citric acid, adipic acid, acidic ion exchange resin, etc.) layered onto or  
 included in the enteric layer, and an outer wall microencapsulated control  
 coating (e.g. polymethacrylic acid ester copolymer or Et cellulose). The  
 multi-walled coated drug has a delayed, gradual, long-term release which  
 takes place in the intestines while the uncoated and/or single-wall coated  
 drug has immediate therapeutic properties upon dissoln. in the stomach.  
 Varying the thickness of the outer control coat affected the release of  
**dextromethorphan**.cntdot.HBr., including a citric acid layer on the  
 inner enteric coating delayed release of the drug.

L10 ANSWER 34 OF 54 CAPLUS COPYRIGHT 2003 ACS on STN  
 AN 1998:723752 CAPLUS  
 DN 129:347319  
 TI Sustained release polymer blend matrix for pharmaceutical application  
 IN Skinner, George William  
 PA Hercules Incorporated, USA  
 SO Eur. Pat. Appl., 15 pp.  
 CODEN: EPXXDW

DT Patent  
 LA English

FAN.CNT 3

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 875245	A2	19981104	EP 1998-107427	19980423
	EP 875245	A3	19990908		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	US 6210710	B1	20010403	US 1997-847842	19970428
	NO 9801893	A	19981029	NO 1998-1893	19980427
PRAI	US 1997-847842	A	19970428		
AB	A pharmaceutical compn. has a blend of at least first and second components and a medicament in a sufficient amt. to be therapeutic where the first component is selected from hydroxypropyl cellulose (HPC), Et cellulose (EC), or derivs. of HPC, EC and hydroxyethyl cellulose (HEC) and the second component is at least one polymer. When HPC is the first component, hydroxypropyl Me cellulose (HPMC), HEC or CM-cellulose will not be the second component and when EC is the first component, HPMC will not be the second component. The medicament can be a variety of drugs or nutritional supplements. The pharmaceutical compn. releases the medicament for a prolonged or sustained period of time and can be formulated into many dosage forms. Formulations of solid oral dosage forms contain phenylpropanolamine and a variety of HPC and CMC or guar.				

L10 ANSWER 45 OF 54 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC. on STN  
DUPLICATE 7  
AN 1994:182906 BIOSIS  
DN PREV199497195906  
TI Mixed ion pair liquid chromatography method for the simultaneous assay of  
**ascorbic acid**, caffeine, chlorpheniramine maleate,  
**dextromethorphan** HBr monohydrate and paracetamol in Frenadol-TM  
sachets.  
AU Thomas, B. R.; Fang, X. G.; Shen, P.; Ghodbane, S. [Reprint author]  
CS Warner-Chilcott Lab., Warner-Lambert Company, 182 Tabor Rd., Morris  
Plains, NJ 07950, USA  
SO Journal of Pharmaceutical and Biomedical Analysis, (1994) Vol. 12, No. 1,  
pp. 85-90.  
CODEN: JPBADA. ISSN: 0731-7085.  
DT Article  
LA English  
ED Entered STN: 26 Apr 1994  
Last Updated on STN: 25 Jun 1994  
AB The five active drug substances and two of the excipients present in  
Frenadol-TM, a cold medication were separated. The active drug components  
**dextromethorphan** HBr monohydrate, **ascorbic acid**,  
caffeine, paracetamol and chlorpheniramine maleate were quantitatively  
assayed by a mixed ion pair LC method. The excipients separated were  
citric acid and maleic acid. The HPLC assay included dual-wavelength  
detection to simultaneously quantify the large concentration of  
paracetamol and the much lower concentration of chlorpheniramine and  
**dextromethorphan**. Both tetrabutylammonium hydrogen sulphate (TBA)  
and pentane sulphonic acid (PSA) were necessary for resolution of the  
seven compounds. The TBA was necessary to lessen peak tailing for  
**dextromethorphan** and chlorpheniramine, to retain **ascorbic**  
acid and to shorten assay time. The pentane sulphonic acid enhanced peak  
shape for **dextromethorphan** and chlorpheniramine. The assay of  
the active drug substances was validated for use in quality control  
applications. Validation studies demonstrated that the procedure was  
accurate, linear, precise, reproducible and rugged. The method conformed  
to both USP and EC validation guidelines.